

Table 23.7  
Cancer of the Prostate (Invasive)

SEER Incidence<sup>a</sup> and U.S. Death<sup>b</sup> Rates, Age-Adjusted and Age-Specific Rates, by Race

	SEER Incidence <sup>a</sup>			U.S. Mortality <sup>b</sup>		
	All Races Males	White Males	Black Males	All Races Males	White Males	Black Males
<u>Age-Adjusted Rates, 2006-2010</u>						
Age at Diagnosis/Death:						
All ages	152.0	144.9	228.5	23.0	21.2	50.9
Under 65	58.8	55.5	101.8	1.8	1.5	4.5
65 and over	796.1	762.5	1,104.3	170.1	157.6	372.1
All ages (IARC world std) <sup>c</sup>	104.6	99.8	162.1	10.0	9.1	22.9
<u>Age-Specific Rates, 2006-2010</u>						
Age at Diagnosis/Death:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	-	-	-	-	-	-
25-29	-	-	-	-	-	-
30-34	0.1	-	-	-	-	-
35-39	1.2	1.0	2.8	0.0	-	-
40-44	10.8	9.2	26.0	0.2	0.2	0.5
45-49	46.5	41.0	101.9	0.9	0.7	2.3
50-54	145.9	135.9	266.6	3.3	2.7	8.1
55-59	327.4	311.5	550.3	9.1	7.7	23.1
60-64	566.7	542.8	922.6	21.0	18.1	53.2
65-69	848.0	816.9	1,266.9	42.8	37.3	107.9
70-74	916.1	885.4	1,251.7	82.8	73.1	206.5
75-79	840.8	802.2	1,083.4	148.7	135.6	344.2
80-84	644.2	604.4	867.9	267.0	249.0	588.8
85+	532.4	503.0	751.4	556.3	529.7	1,094.4

<sup>a</sup> SEER 18 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

<sup>b</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

<sup>c</sup> Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

- Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.